Serial No.: 09/222,460 Filed: December 29, 1998

LISTING OF CLAIMS

- 1. (currently amended) A method for the treatment of metanephric tissue for transplantation into a recipient comprising:
- <u>a.</u> contacting said metanephric tissue, *in vitro*, with a growth factor-containing composition comprising one or more growth factors for metanephric development <u>vascular endothelial growth factor;</u> and
 - b. transplanting said metanephric tissue into said recipient.
 - 2-3. (canceled)
- 4. (previously presented) The method of claim 1 wherein said metanephric tissue is contacted with said growth factor-containing composition for less than 8 hours.
- 5. (previously presented) The method of claim 1 wherein said metanephric tissue is contacted with said growth factor-containing composition for less than 2 hours.
 - 6. (canceled)
- 7. (currently amended) A method for the treatment of metanephric tissue transplanted into a recipient comprising contacting said transplanted metanephric tissue with a growth factor-containing composition comprising one or more growth factors for metanephric development vascular endothelial growth factor, wherein said growth factor-containing composition is administered to said transplanted metanephric tissue at the time a ureteroureterostomy is performed.
- 8. (previously presented) The method of claim 22 wherein said growth factor-containing composition is administered to said metanephric tissue by an osmotic pump.

Serial No.: 09/222,460 Filed: December 29, 1998

9. (currently amended) The method of claim 22 wherein said growth factor-containing composition is administered to said recipient in a manner such that said one or more growth factors for metanephric development are vascular endothelial growth factor is present in said recipient's blood that circulates through said metanephric tissue.

10-21. (canceled)

22. (currently amended) A method for the treatment of metanephric tissue comprising contacting said metanephric tissue, *in vivo*, with a growth factor-containing composition comprising one or more growth factors for metanephric development vascular endothelial growth factor at the time of or after being said metanephric tissue is transplanted into said recipient.

23-24 (canceled)

- 25. (currently amended) The method of claims 1, 7, or 22, 26, 29, or 30 wherein said contacted metanephric tissue develops into a functional chimeric kidney in said recipient and wherein the glomeruli of said functional chimeric kidney are vascularized primarily by said recipient and are able to filter plasma.
- 26. (new) A method for the treatment of metanephric tissue for transplantation into a recipient comprising:
- a. contacting said metanephric tissue, *in vitro*, with a growth factor-containing composition comprising vitamin A; and
 - b. transplanting said metanephric tissue into said recipient.

Serial No.: 09/222,460 Filed: December 29, 1998

27. (new) The method of claim 26 wherein said metanephric tissue is contacted with said growth factor-containing composition for less than 8 hours.

- 28. (new) The method of claim 26 wherein said metanephric tissue is contacted with said growth factor-containing composition for less than 2 hours.
- 29. (new) A method for the treatment of metanephric tissue transplanted into a recipient comprising contacting said transplanted metanephric tissue with a growth factor-containing composition comprising vitamin A, wherein said growth factor-containing composition is administered to said transplanted metanephric tissue at the time a ureteroureterostomy is performed.
- 30. (new) A method for the treatment of metanephric tissue comprising contacting said metanephric tissue, *in vivo*, with a growth factor-containing composition comprising vitamin A at the time of or after said metanephric tissue is transplanted into said recipient.
- 31. (new) The method of claim 30 wherein said growth factor-containing composition is administered to said metanephric tissue by an osmotic pump.
- 32. (new) The method of claim 30 wherein said growth factor-containing composition is administered to said recipient in a manner such that said vitamin A is present in said recipient's blood that circulates through said metanephric tissue.